L5 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2008 ACS on STN

AN 2004:1023502 CAPLUS Full-text https://chemport.fiz-

karlsruhe.de/cgi-

bin/ex sdcgi?yyUE0ijzB0 PDnf0qaWiU wI3RGzGCmJbFx0sibuzToprnspWdxrkGMudGGpEcmcY4ShdeVV3pXiVVtQxZOBe az7FX7lcmsSp7D0SWKKsdQBY34Wilr6ZBRPRKQVgFkSyzvkewpUEPGK 5D qC6wWzxBYp6YtjDH55f6WsUc_5VxBFGZB8p7e70taR2A2S5yH>DN 141:410956

TI A preparation of pyridothienopyrimidine derivatives, useful as herbicidal

antidotes
IN Vasilin, V. K.; Osipova, A. A.; Kaigorodova, E. A.; Nen'ko, N. I.;
Krapivin, G. D.; Isakova, L. I.; Strelkov, V. D.

PA Kubanskii Gosudarstvennyi Tekhnologicheskii Universitet, Russia

SO Russ., No pp. given

CODEN: RUXXE7

DT Patent

LA Russian

PATENT N	0.	KIND	DATE	APPLICATION NO.	DATE
PI RU 22410	02	C1	20041127	RU 2003-123516	
20030724 <					
PRAI RU 2003-	123516		20030724		
GI					

AB The invention relates to a preparation of pyridothienopyrimidine derivs. of formula I (wherein: R is 2-(5-bromo-furyl), 2-furyl, and cyclohexyl], useful as herbicidal antidotes of 2,4-dichlorophenoxyacetic acid. For instance, pyridothienopyrimidine derivative II was prepared via heterocyclization of thienopyridine derivative III and 5-bromo-2-furfural with a yield of 83%. In the presence of compound II, toxic effect of herbicidal 2,4-dichlorophenoxyacetic acid decreased (at the concentration of 10-2% sunflower root length increased by 30%).